MISSOURI BO

PETER H

GEO

BOTA



Sinc

page

onlin

nu

27

Wight ic. 1371

1371 C. Arabica, (sp. Linn. acuta) each fl. acute or pedicellate,
calyx ½ lin. long glaby. ovoid and the base of corolla
calcs scarcely conspicuous or wanting
Seinde, Stocks, on Arnottas observations etc.

1372 C. hyalina / Dunals, 3 flowers fl. pedicellate, corolla 4-5
deft. hyalin et Falconeottah on Amer. observ.
Islands at Ceylon. hyal. & subc.

1373 C. Thwaitesii fl. pedicellate acute or subacute
minute pruinose-glaucous, insects — the throat
Ceylon on Vinca rosea — " fascicles of flowers pedicels
or glaucous — calyx & corolla hairy & outside
or linear " Thwaites

Bentham : Fl. Hong : 1871 p. 219

C. Griff / Ophthyrsa v. Cuscuta minor

Ophthyrsa vulgaris ~~ophthyrsa~~ v. ex Thyrsa viridi
ablatam

2, Ophthyrsa capitulis flavis minutis
pedunculatis plantis quae in silvarum parvis
foliis Thyrsoidea hirsutis (et Cuscuta
an sine alia pleraque!) — utrin & fructus

Wight Ill. p. 230.

C. major.

Griff. tab. 61 Wag 243 notulae

Arab. Gujarat / tab 61 in Lawton of a Convolvulus

2 diff Spang— tab 166 simplex
Griffith Jan March
tab 168. same.
apparently one plant. the simplex the
state — only a Gentian plant.

Flora Ind. p. 220. —

C. Langsd. fl. pubiscell. ? gen end
C. Griffy. fl. zeaid, fl. ambelli,
a. latiorifl, Hegn, Griff Lapponici
Spreng. = rubioides?

p. 231 C. Gunnisoni Jan Lessing a Orion Davilion
fl. pubescentia, tubo cor. subtilissime sericeo-pubescente,
limbo subbrevius, filamentis basi dilatatis
intus ciliatis, ovario glabro, stylis ovario
brevioribus, stigm capit.

fl. rubra glaucescenti, capsulis
breviss vel longiss pedicellat, cal 4 part,
lob ovat. Corolla subtilissime barbata
obtus ovatis calyce paulo longioribus, ad
basin cap usque 4 fida, lacinia 4 falcata
lanceolatis; stam incl. ; ng. nullis

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

5120

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

28

/ Cuscuta Europaea β. racemosa Gren & Godr Fl. de France, II, 504

caulibus nullis, capsula ovata
the stipes itself: capsula obpyriforme a summit
attenuata. Flores sessiles

C. Epithymum ibid fl. sessiles petites (moitié
moindres que celles du C. Europaea); cal.
lob. ovat acuminat, apice patulis; corol.
triangulat; squam magnis tubum claudent
styles exsertis, dans stamens suspentibus
a Cotinus sec, commun dans le midi &.

C. Trifolii cal lob. adpressis; corol. angulatibus
longioribus, sq. angulatoribus tubum totum non
claudentibus, styles demersatis, nunquam stamens
suspentibus.

C. alba ibid lobis et squamis subobtusis v.
obtusis — flowers smaller & more compact
glomerules smaller — more common than this
than Epithymum; — more common than this
dans toute la région méditerranéenne.

" * is this my C. mediterranea? or this
and Epith united?

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

5121

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

29/ Bull. de la Société botanique de France I. p. 295 1854

Gramnitis Loureiro Flor Cochinchinensis

Des Moulins

Cal. perianth. inferum 5 fidum, laciniis antici colorati
ramosis

Cor. campanulat membranea 1 petala, tubo globoso
calyci aequali, limbo brevi 5-fido, laciniis
rotundis patentibus.

Stam. filamenta 5 tubulata corollae aequalia,
tubo insistentia, antheris ovatis incumbentes,

Pist. germ. subrotundum stylo 2 aequali,
stamina. stigmata compresso-rotunda

Pericarp. bacca membranacea subrotunda 4-loba
1 locularis 4 sperma

Sem. subrotunda.

Nom (γραμμή, linea) a forma
plantae linearis.

Gramnitis aphylla (ou low plants in rather
places, in Cochinchina)

Said to be = Banks herbarium, the Conservator of
Such is Rob. Brown,

Des moulins:

Gramnitis suaveolens (no Comments) Swartz
" acuminata (") Linn
G. chrysoroma (") Wallich
G. arabica (") Fresen

5122

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

Grammitis

Archives de Flore, par F. Schultz / mars 1855

p. 81. Mon frère vient de m'écrire, que dans un discours prononcé dans la séance de Manheim il a réuni le C. pumilum au genre Grammitis de Lowrieo, mais je ne pense pas qu'il appartienne à ce genre

p. 145 : Desmoulins dans le bulletin publié en janvier 1855 adopte le nom de Grammitis



mais mon frère dans son discours prononcé
le 15 Nov. 1846 à reçu dyx (= cinquante)
de ces Capitalistes - (Capitalistes spec-
ulateurs) un jeune Gramen !

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

5/23



Gussone Florae Siculae Synopsis
vol I. Neapoli 1842

pag 290. C. alba Prsl 87.. Guss. pr. I. p. 291
flor glomerat sessil 5-fid, laciis ovatis
calycinisque obtusis - ng. hypostamii deutzolat.
stige subcapitat
Caules ramis rubentes, bracteae numeri-
ores et flores minores quam in sequente
(C. Epithymum) cujus forsan var. nequis varietas.

C. Epithymum — flor albi in glomerulis
multifloris dense congesti, calycis basi virentes
laciniis ramulis erectis, corollae albae laciniis
sub anthesi erecto-patentibus tubo aequi-
longis, autumno hiemeque vertue

vol. II pat 2. (1844) p. 887. C. micrantha Ten
[Etna mnd be the Palermo-alba]: flor sessil
4 fid. laciis cal. ovall obtus! (v.s.) in
aridis aprilibus passim Neapolitanis, supra
l'Astrone e Grotto delle Capre.
2-5-flores in glomerule
p. 888. C. tubulata Ten flor. glom. sessil 5-aculeis,
laciis ovoll calycis laureolato-tubulatis!
(v.s.) squamis hypostam. deutzolatis, stig
bicanthus;
In aridis maritimis vel submontanis apricis
passim Zizyphi Lotis, Calothamni, Smilacis
et. — Palermo, Monte Peregrino etc.
flores minores quam in C. alba, bracteis in glom.
multifloros densis congestis; calycis basi rubentibus corollis
subbrevioribus

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

5124

Missouri Botanical Garden
George Engelmann Papers

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
Missouri Botanical Garden

Tenore, Flor. Neap.
Neap. 1824-29

st

small white heads

Tenore tab CCXX fig 3

Cussonia planiflora III. pag 250.

(published 1824-29)

Caulibus capillaribus, flor. glomeratis sessilibus 5 fidis corollis subrotatis, laciniis ovatis obtusis patentissimis staminibus exertis basi squama perexigua bifida munitis; stylis filiformibus, divaricatis; stigmatibus conformibus obtusiusculis.

capolini del diametro di una linea ad una linea e mezza

Alla Plantago lanceolata [Napoli] fiorisce in Maggio.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

5125



MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

39

Jenore flor. hesp. III. p. 249

C. alba Besl 87 Jusne i. p. 291 floribus glomeratis subpedunculatis (nec sessilibus) foliis ovatis lobis ~~ovatis~~ erectis subrotundis obtusis, staminibus basi squama denticulata munitis stigmatibus capitatis.

heads 3 lines diameter.

C. Cynthymeus ovatis lobis patentibus acuminatis staminibus inclusis basi squama ovata munitis.

vol V. p. 335.

C. Cardunculus, Aloysio, in Humulus & Annalis.



5126

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

34

Diagnoses plantarum novarum praesertim orientalium
et
Series secunda nro. 3, E. Boissier
Lips. 1856

pag 126. Cistineae.

C. globulosa Boiss & Reut. in valle alpina Bozdagh.
notiert Tmoli occidentali, ad Genistas, Balansa 1854 Nr 413

C. macranthera Heldr & Sartori Jagyd. & Parnass. ad
notiert Daphnen buxifol & Sideritid.

p. 128 C. Cassiopes [Calliopes ?] Heldr & Sartori ad Parnassi
notiert regione alpina ~~ad~~ Daphnen buxifol.

p. 129. C. elegans Boiss & Balansa, inter Ciliciam
notiert et Caesaream, in regione alpina montis Masmenen
Dagh, ad Convolv. compact. et Genistas.

p. 129 C. ciliaris Hohenack = Kotschy N. Assyr. (for the first
time descr. here.

p. 130. C. oxypetala Boiss = C. Arabica Wight. Ic. tab 1371
non Presen. In Stirade Ins. or. ad. Triantheman
Stocks leg.

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

5127

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

25

Notes fr. Acom. & Shortt
Edit. Linn. Bot. VI. 1820.

Cassuta pag 199

C. corymbosa a.v.D. 4partel, with the scales, a medio sativa
C. floribunda K. corymbis pedunculatis, carinis obtusis
Corollis globosis, calyce depte superiibus, carinis
acutis reflexis genitalibus longe exsertis, segm
reflexis. C. Gopsycamensis valde affinis.
Nov. Hisp. prope montem Ilt Istla
C. asperrima Th. pedunculis unifloris, capillaribus
calyce 5 fidis brevissimis. pilosa, carinis acutis
flavis minores quam C. acuminata
C. monogyna Vahl. Orientis. Jacquem. Gelliz
prope Sangarium, et Caucaso.
Vahl Symb. II. 92
C. longistylosa Kn. staminalis
C. chinensis Lam. fl. staminalibus; calyce angulosa
longitudine Corollae; corolla C. carinis t. roseis
alatum 5 partum. stigm. subcapitata
C. reflexa Roxb. fl. roseis, bracteis pedicellatis; calycis
foliolis ovatis, corolla tubo cylindr.; segmentis
barbatis carinatis tubi inaequalis stylo vix ullius
stigmata exserta acuminata. Coromandel
C. australis Br. ad Port. Jackson. stig. exsertis
segmentis. Latitudo fi lamentis subulatis; calycis
segmenta profunde quinquefid. Caes.
C. racemosa Roxb. fl. roseis, bracteis pedicellatis; calycis
foliolis ovatis, corolla tubo cylindr.; segmentis
barbatis carinatis tubi inaequalis stylo vix ullius
stigmata exserta acuminata. Coromandel
Cassia tomentosa (Grammica asperella) et. Nov Holl tropica

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

C. hyalina Roth; pedunculo tripluri corolla calyce majori,
laciniis ~~ovatis~~ lanceolatis Roth nov. pl. spec
Mass
ind. cum v. Hegne
capillaris, corolla 4-fida calyce dimidio majore
stamina squamis (!!) imposita ori corollae affixis
C. fascicularis
C. Gmelinii Willd. Spec. nov. Vol 1. Addg. Willd
C. cymosa Willd. an umbellata (an! ex Ind
C. hoppeana ic

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

5128

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN

medici Surinamensis

J. G. Schilling v. de Lepra

Commentationes

Lugd. Batav. 1778.

pag. 200. Cancrata Surinamensis

(pag. 60: In tractu Surinamensi frequentissima est Cancrata spuria, quae nullas arbores, vivam in modum infestat, obvolvens sarmentis suis, quae valde tenacia sunt, omnes pedetentim ramos, tandem universam ~~magnam~~ quodam obtegens.

This Cancrata (no bad odor is mentioned
— vide C. graveolens HBK! perhaps is the same)
is macerated till it ferments a day
or two begins to stink, & the milky juice
made by ~~cont~~ agitation, working &
kneading into an ointment used against
Lepra or Elephantiasis.

pag. 200. ~~Floribus~~
Cancrata Surinamensis
floribus fasciculos undique pandit
mire prolifera.
Calyx 4 phyllus, 5 fidus — Lacinia
filamenta 10 non distincta sed ad
angulos corollae adnata. Stylus 2 stigmata
tuberculata, capsula tuberculata

5129

Tab II. a figure of the plant.

fig 2

calyx & corolla

fig 3



top view of Corolla

fig. 4



fig 5.

MISSOURI BOTANICAL GARDEN
GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10
cm
copyright reserved
MISSOURI BOTANICAL GARDEN